

Mortality

| Hospital Quality Measures | What Is This? Why Is It Important? |
|--|--|
| 30-Day Death Rate for Chronic Obstructive Pulmonary Disease (COPD) Patients | <ul style="list-style-type: none"> ◆ The death rates show whether patients died within 30 days of being hospitalized for a specific condition or having had surgery. ◆ Death rates provide information about important aspects of hospital care that affect patients' outcomes – like prevention of and response to complications, emphasis on patient safety, and the timeliness of care. ◆ Lower Percentages are Better. |
| 30-Day Death Rate for Heart Attack Patients | |
| 30-Day Death Rate for Heart Failure Patients | |
| 30-Day Death Rate for Pneumonia Patients | |
| 30-Day Death Rate for Stroke Patients | |
| 30-Day Death Rate for Coronary Artery Bypass Graft (CABG) Surgery Patients | |
| Deaths (per 1,000) Among Patients with Serious Treatable Complications After Surgery | <ul style="list-style-type: none"> ◆ This measure refers to surgical patients who died after developing serious complications that could have been treated. ◆ There is always a risk of serious complication during or after surgery. However, hospitals with trained, well organized, and efficient staff identify these complications quickly and treat them aggressively. ◆ Some deaths may be unavoidable. Higher death rates from complications may be a sign that patients were not watched closely after surgery or that effective action was not taken. ◆ Lower Numbers are Better. |

Mortality

| Hospital Quality Measures | * Indicator | Performance Period | Dearborn | Farmington Hills | Grosse Pointe | Royal Oak | Taylor | Trenton | Troy | Wayne | National Benchmarks |
|---|-------------|-----------------------|----------|------------------|---------------|-----------|--------|---------|--------|-------|---------------------|
| 30-Day Death Rate for Chronic Obstructive Pulmonary Disease (COPD) Patients | ↓ | July 2013 – June 2016 | 5.8% | 7.7% | 6.7% | 6.1% | 8.9% | 7.7% | 7.0% | 9.8% | 8.0% |
| 30-Day Death Rate for Heart Attack Patients | ↓ | July 2013 – June 2016 | 11.0% | 12.9% | 10.8% | 11.4% | 13.4% | 12.7% | 10.6% | 13.2% | 13.6% |
| 30-Day Death Rate for Heart Failure Patients | ↓ | July 2013 – June 2016 | 9.6% | 10.0% | 8.3% | 8.3% | 11.0% | 9.6% | 8.7% | 10.9% | 11.9% |
| 30-Day Death Rate for Pneumonia Patients | ↓ | July 2013 – June 2016 | 13.5% | 15.8% | 11.7% | 11.8% | 13.7% | 14.3% | 12.5% | 15.7% | 15.9% |
| 30-Day Death Rate for Stroke Patients | ↓ | July 2013 – June 2016 | 14.4% | 13.6% | 13.3% | 12.1% | 13.1% | 16.2% | 12.2% | 14.1% | 14.6% |
| 30-Day Death Rate for Coronary Artery Bypass Graft (CABG) Surgery Patients ¹ | ↓ | July 2013 – June 2016 | 2.7% | N/A | N/A | 3.0% | N/A | N/A | 3.1% | N/A | 3.2% |
| Deaths (per 1,000) Among Patients with Serious Treatable Complications After Surgery ² | ↓ | July 2013 – June 2015 | 139.53 | 143.01 | 127.15 | 133.34 | N/A | 128.54 | 126.67 | N/A | 139.05 |

Footnote Legend:

¹ N/A: CABG Surgery Not Performed at this Hospital.

² N/A: Number of Cases/Patients is Too Few to Report at this Hospital.

| * Indicator | |
|-------------|--|
| ↑ | Higher Values Signify Better Performance |
| ↓ | Lower Values Signify Better Performance |

| Colors Indicate Beaumont Performance (Based on Confidence Intervals) Against National Performance | | | |
|--|------------------|-------------------|--------------------|
| Better Performance | Same Performance | Worse Performance | N/A Not Applicable |

| We Report All Our Data to CMS and CMS Reports Our Data through Medicare.gov | |
|---|--|
| Learn More At: | beaumont.org/quality |
| | medicare.gov/hospitalcompare |